

White, W. (2011). Smoking and addiction recovery. For people in recovery.
Posted at www.williamwhitepapers.org.

Smoking and Addiction Recovery: For People in Recovery

William L. White

For decades, people in recovery from addictions to other drugs have had their lives cut short by tobacco-related diseases. These dear friends, patients, and colleagues died from nicotine addiction, but it could also be said they died from blindness—the failure to see nicotine as an addictive drug and the failure to embrace the need for smoking cessation within our understanding of *recovery*. For years, such casualties could be written off to the lack of knowledge about smoking and health and the lack of knowledge about the effects of smoking on recovery from other addictions. That simply is no longer the case. Science has weighed in on these issues, and the findings are excruciatingly clear. Here are the facts.

Nicotine is an addictive drug.¹

- **Genetic Influences**: Genetic vulnerability plays a role in who starts smoking, who develops nicotine dependence, and who succeeds with particular approaches to quitting.² Families with histories of alcohol addiction have high rates of nicotine addiction—even among family members who are not dependent on alcohol.³
- **Difficulty Quitting**: Most smokers regret starting to smoke (90%), want to quit and have made a prior attempt to quit (70%),⁴ or are currently trying or planning to quit (40%).⁵ A higher percentage of people successfully recover from alcohol and other substance use disorders than successfully quit smoking.⁶
- **Cycles of Use and Abstinence**: Like other addictions, recovery from nicotine addiction involves cycles of abstinence, lapse, relapse, remorse, remotivation, and new abstinence attempts.⁷ Those who attempt to quit smoking report an average of 8-11 quit attempts before succeeding, with most relapsing within the first eight days following their last cigarette.⁸

Nicotine has been, until recently, a celebrated drug.

- Hidden Addiction: The problem with drugs that have been so celebrated—culturally praised, commercially promoted, and politically protected—is that we have not defined them as *drugs*, have not regulated them as we have other drugs, have not defined their compulsive use as *addiction*, and as a result, have only recently begun to fully understand their relationship to disease and premature death.⁹

Nicotine addiction is declining, but remains a significant public health problem.

- Current Progress: Although the rate of smoking for adults in the US has declined from its peak of 45% in 1954 to 20.6% in 2009, 46.6 million Americans are currently addicted to nicotine.¹⁰

The medical consequences of nicotine addiction remain a leading cause of death in the United States.

- Disease and Death from Nicotine Addiction: Tobacco use accounts for more sickness and disease than the combined use of alcohol and other drugs.¹¹ Medical disorders related to nicotine addiction cause more than 430,000 deaths each year in the United States;¹² cigarette smoking is the leading preventable cause of death in the United States.¹³ One-third to one-half of lifelong smokers will die of smoking-related illnesses.¹⁴
- Added Risk from Co-addiction: Combining nicotine addiction with other drug addictions amplifies the health risk of both addictions.¹⁵ For example, the risk of laryngeal cancer for those who are both heavy drinkers **and** heavy smokers is four times greater than for those who are either heavy drinkers or heavy smokers.¹⁶

Co-addiction is the norm.

- Smoking and Drinking: Nicotine use increases alcohol consumption; alcohol consumption increases nicotine consumption and the rewarding effects of nicotine.¹⁷ Alcohol dependent smokers are more severely dependent upon nicotine,¹⁸ are less likely to stop smoking,¹⁹ and as a result, bear a greater burden of nicotine-related diseases and deaths.
- Smoking and Psychiatric Illness: Overall, 40% of smokers in the US have a substance use or psychiatric disorder.²⁰
- Smokers in Addiction Treatment: Between 70-80% of people entering addiction treatment smoke—nearly 4 times the rate for all adults.²¹

- Co-addiction and Prognosis for Quitting: Smoking cessation rates are extremely low for alcohol and drug dependent persons not in recovery from these disorders; recovery from other alcohol/drug dependence may be a precondition for smoking cessation for most addicted smokers.²²

Nicotine addiction (smoking) is a major cause of death among people in recovery from other addictions.

- Causes of Death Following Addiction Treatment: People treated for alcohol or drug dependence are more likely to subsequently die from smoking-related diseases than from alcohol or drugs other than nicotine.²³

Many of the pioneers of twentieth century addiction treatment and recovery mutual aid societies died of smoking-related disorders.

- Bill Wilson (emphysema) and Dr. Robert Holbrook Smith (cancer), co-founders of Alcoholics Anonymous;
- Mrs. Marty Mann (cancer), founder, National Council on Alcoholism and Drug Dependence;
- Danny C. (cancer) and Jimmy K. (emphysema and cancer), key figures in the founding of Narcotics Anonymous;
- Charles Dederich (cardiovascular disease), founder of Synanon;
- Dr. Marie Nyswander (cancer), co-developer of methadone maintenance; and
- Senator/Governor Harold Hughes (emphysema), sponsor of landmark alcoholism treatment legislation and founder of the Society of Americans for Recovery.²⁴

Quitting smoking reduces the risk of relapse and increases one's odds of sustained recovery from alcohol and other drug addictions.²⁵

- Risk of Continued Smoking: Smoking cessation in early recovery does not increase craving for or urges to drink alcohol.²⁶ Continued smoking following treatment for alcohol and drug dependence is a risk factor for resumption of alcohol and other drug use.²⁷
- Improved Recovery Rates: Quitting smoking in tandem with efforts to cease alcohol and other drug use does not threaten such recovery efforts and in fact, increases one's chances of recovery.²⁸ Smoking cessation improves recovery rates of other addictions.²⁹

There are medications and counseling available that can increase the odds of permanent smoking cessation.

- Value of Medication: Medications can double the chances of successfully quitting,³⁰ and nicotine replacement medications may be particularly important for smokers with a history of alcohol and/or drug problems.³¹
- Value of Medication and Counseling: Smokers who combine medications and counseling have the highest success rates.³²
- Value of Physician Guidance: Smokers with co-occurring addictions are five times more likely to successfully quit smoking if they are counseled by a primary care physician.³³

People in recovery from addiction to alcohol and other drugs are incorporating recovery from nicotine addiction within their personal understanding of recovery. This is revealed by:

- the rise of smoking cessation with length of abstinence from alcohol and other drugs,³⁴
- the growth of non-smoking meetings within recovery fellowships,
- some people in recovery changing their sobriety/clean dates to reflect the date they stopped all addictive drug use—including nicotine use, and
- addiction professionals broadening their understanding of “recovery” to encompass smoking cessation.³⁵

The health benefits of smoking cessation include increased life expectancy; reduced risk of heart disease, heart attacks, strokes, and cancer;³⁶ as well as a more rapid process of brain recovery from addiction.³⁷

10 Suggestions for People in Recovery Who Wish to Quit Smoking

1. Expect success in quitting. Full and permanent recovery from nicotine addiction is possible for people recovering from other addictions. More than 48 million Americans have successfully quit smoking,³⁸ and persons with a history of other drug addictions can also successfully quit.³⁹ You can do this, but it will require sustained effort. Do not be discouraged. Many of us had to keep quitting until we quit completely and permanently.

2. Educate yourself. Do an internet search on “smoking cessation” and get more facts on the effects of smoking cessation on your health.

3. Prepare yourself by making a quit plan that includes these steps.

- Focus. (Write down all the reasons you want to quit smoking.)
- Set a quit date.
- Announce your quit date to others.
- Take preliminary steps, e.g., start decreasing the number of cigarettes you smoke each day; create a smoking diary by recording when you smoke, noting times, places, people, situations; experiment by eliminating a few of those “special” cigarettes (during phone calls, work breaks; while driving; after a meal; after sex—you know the ones I am talking about).
- Talk to your partner. (If you live with a smoker, explore whether you want to quit together. You will be five times more likely to give up smoking if he or she also quits.⁴⁰)
- Talk to your doctor about your decision and the options for medications that may be of help to you.
- Line up additional sources of support. Line up lots of support and be prepared to combine multiple methods of support, such as self-instructional materials, medications, counseling, and mutual support (e.g., Nicotine Anonymous). Although the majority of people who quit smoking do so without assistance,⁴¹ the use of self-help groups, medications, counseling, and other aids to support smoking cessation have increased dramatically over the past three decades⁴² and will increase your chances of being successful.

4. Apply principles and practices that have helped you recover from other addictions.⁴³ Although persons with additional drug addictions tend to have more severe nicotine dependencies, people in recovery from such addictions may be able to mobilize skills and supports that helped them recover to aid in their efforts to quit smoking.⁴⁴

5. Talk with other people in recovery who have stopped smoking. Ask them to share their experiences and suggestions related to quitting.

6. Identify your most powerful smoking triggers. Use your smoking diary to identify the people, places, things, times, situations, and feelings most associated with smoking and minimize contact with these, particularly during the first 30 days of not smoking.

7. Minimize and manage cravings by doing the following:

- Closely adhere to medications if you are using them to suppress cravings.
- Use thought stopping (an internal shout “Stop!”) that can be further enhanced by wearing a rubber band on your wrist and snapping it every time you think of having a cigarette. (I know this sounds silly, but it really works in extinguishing cravings.)
- Combine thought stopping with thought substitution—have a single thought (mantra) or image you can bring to mind as an alternative to thinking about smoking that strengthens your resolve to quit.
- When you experience a craving, change your physical environment for a few minutes.
- Increase your water intake—this may help with nicotine detoxification and carrying a bottle of water and drinking from it gives you an oral substitute and an alternative activity for your hands.
- Substitute oral activity for smoking during the early weeks of not smoking, e.g., sugarless gum, mints, toothpicks, straws, healthy snacks (e.g., carrot sticks).
- Increase your exercise.
- Carry an icon with you that you can draw on for strength, e.g., photo of a loved one, photo of someone who died of nicotine addiction, a worry stone, a recovery chip.
- Structure your time to remain in non-smoking environments (e.g., non-smoking recovery support meetings), particularly for the first month.
- Use other techniques, including breathing exercises, visualization, and daily sharing with a quit mentor.

8. Anticipate the need to relearn how to do some daily tasks without smoking. (Smoking had become so integrated into my writing rituals for so many years, I had to quite literally relearn how to write when I quit smoking.)

9. Plan rewards for yourself to celebrate key recovery milestones, e.g., first day, first week, first month.

10. For other great suggestions, check out <http://www.becomeanex.org/> and join the Ex Community.

Dedication: This series of papers is dedicated to Charlie B.

Acknowledgement: Many of the findings in this review were abstracted from the published studies and accompanying literature citations of Hannah K. Knudsen, PhD (Department of Behavioral Science and Center on Drug and Alcohol Research, University of Kentucky) and her colleagues. These studies, which were supported by the National Institute on Drug Abuse, have significantly advanced our understanding of the barriers to implementing comprehensive smoking cessation supports within addiction treatment programs in the United States. Her contributions in leading this research are gratefully acknowledged. I'm also indebted to the research that has been collected and synthesized by the American Legacy Foundation (see <http://www.legacyforhealth.org/>) and to David B. Abrams, PhD, for his many helpful suggestions. Development of this series of papers was supported by the Philadelphia Department of Behavioral Health and Intellectual disAbility Services.

About the Author: William White is a Senior Research Consultant at Chestnut Health Systems and has served as a volunteer for Faces and Voices of Recovery since its inception in 2001. He has worked full-time in the addictions field since 1969 and quit a two-decades-long, two-pack-a-day addiction to nicotine on January 1, 1988.

¹ Bonnie, R. J., Stratton, K., & Wallace, R. B. (Eds., 2007). *Ending the tobacco problem: A blueprint for the nation*. Washington, D.C.: Institute of Medicine.

² Bonnie, R. J., Stratton, K., & Wallace, R. B. (Eds., 2007). *Ending the tobacco problem: A blueprint for the nation*. Washington, D.C.: Institute of Medicine.

³ Cuijpers, P., & Smit, F. (2002). Nicotine dependence and regular nicotine use in adult children of alcoholics. *Addiction Research & Theory, 10*, 69-81.

⁴ Hughes, J. R., Marcy, T. W., & Naud, S. (2008). Interest in treatments to stop smoking. *Journal of Substance Abuse Treatment, 36*, 18-24.

⁵ Bonnie, R. J., Stratton, K., & Wallace, R. B. (Eds., 2007). *Ending the tobacco problem: A blueprint for the nation*. Washington, D.C.: Institute of Medicine, p. 88.

⁶ 51% lifetime remission for cigarette smoking. Centers for Disease Control and Prevention. (2009). Cigarette smoking among adults and trends in smoking cessation—United States, 2008. *MMWR Morbidity and Mortality Weekly Report, 58*(44), 1227-1232. 45-80% for alcohol and drug use disorders. Compton, W. M., Thomas, Y. F., Stinson, F. S., & Grant, B. F. (2007). Prevalence, correlates, disability, and comorbidity of DSM-IV drug abuse and dependence in the United States. *Archives of General Psychiatry, 64*, 566-576. Dawson, D. A. (1996). Correlates of past-year status among treated and untreated persons with former alcohol dependence: United States, 1992. *Alcoholism: Clinical and Experimental Research, 20*(4), 771-779. Hasin, D. S., Stinson,

- F. S., Ogburn, E., & Grant, B. F. (2007). Prevalence, correlates, disability and comorbidity of DSM-IV alcohol abuse and dependence in the United States: Results from the National Epidemiologic Survey on Alcohol and related conditions. *Archives of General Psychiatry*, 64(7), 830-842. Kessler, R., McGonagle, K., Zhao, S., Nelson, C., Hughes, M., Eshelman, S.,...Kendler, K. (1994). Lifetime and 12-month prevalence of DSM-II-R psychiatric disorders in the United States. *Archives of General Psychiatry*, 51, 8-19.
- ⁷ Bonnie, R. J., Stratton, K., & Wallace, R. B. (Eds., 2007). *Ending the tobacco problem: A blueprint for the nation*. Washington, D.C.: Institute of Medicine.
- ⁸ U.S. Department of Health and Human Services (2001). *Women and smoking: A report of the Surgeon General*. Rockville, MD: U.S. Department of Health and Human Services, Center for Disease Prevention and Health Promotion, Office on Smoking and Health. Hughes, J. R., Keely, J., & Naud, S. (2004). Shape of the relapse curve and long-term abstinence among untreated smokers. *Addiction*, 99(1), 29-38.
- ⁹ White, W. (1996). *Pathways from the culture of addiction to the culture of recovery*. Center City, MN: Hazelden.
- ¹⁰ Centers for Disease Control and Prevention. (2010). Vital signs: Cigarette smoking among adults aged >18 years—United States, 2009. *Morbidity and Mortality Weekly Report*, 59(35), 1135-1140. Saad, L. (2008). *U.S. smoking rate still coming down*. Retrieved from <http://www.gallup.com/poll/109048/us-smoking-rate-still-coming-down.aspx>?
- ¹¹ U.S. Department of Health and Human Services. (2000). *Healthy People 2010: Understanding and improving health*. Washington D.C., Author.
- ¹² CDC. (2008). Smoking-Attributable mortality, years of potential life lost, and productivity losses --- United States, 2000--2004. *Morbidity and Mortality Weekly Report*, 57(45), 1226-1228. Kalman, D., Kim, S., DiGirolam, G. Smelson, D., & Ziedonis, Z. (2010). Addressing tobacco use disorder in smokers in early remission from alcohol dependence. *Clinical Psychology Review*, 30(1), 12-24.
- ¹³ CDC. (2002). Annual smoking-attributable mortality, years of potential life lost and economic costs United States, 1995-1999. *Morbidity and Mortality Weekly Report*, 51, 300-303.
- ¹⁴ U.S. Department of Health and Human Services. (2004). *The health consequences of smoking: A report of the Surgeon General*. Atlanta, GA: U.S. Department of Health and Human Services, Center for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health. p 873.
- ¹⁵ Kalman, D., Kim, S., DiGirolam, G. Smelson, D., & Ziedonis, Z. (2010). Addressing tobacco use disorder in smokers in early remission from alcohol dependence. *Clinical Psychology Review*, 30(1), 12-24.
- ¹⁶ Gulliver, S.B., Kamholz, B.W., & Helstron, A.W. (2006). Smoking cessation and alcohol abstinence: What do the data tell us? *Alcohol Research & Health*, 29(3), 208-212.
- ¹⁷ Kalman, D., Kim, S., DiGirolam, G. Smelson, D., & Ziedonis, Z. (2010). Addressing tobacco use disorder in smokers in early remission from alcohol dependence. *Clinical Psychology Review*, 30(1), 12-24.
- ¹⁸ Bien, T. H., & Burge, R. (1990). Smoking and drinking : A review of the literature. *International Journal of the Addictions*, 25, 1429-1454. Kalman, D., Kim, S., DiGirolam, G. Smelson, D., & Ziedonis, Z. (2010). Addressing tobacco use disorder in smokers in early remission from alcohol dependence. *Clinical Psychology Review*, 30(1), 12-24. York, J. L., & Hirsch, J. A. (1995). Drinking patterns and health status in smoking and nonsmoking alcoholics. *Alcoholism: Clinical and Experimental Research*, 19(3), 666-673. Burling, A. S., & Burling, T. A. (2003). A comparison of self-report measures of nicotine dependence among male drug/alcohol dependent cigarette smokers. *Nicotine and Tobacco Research*, 5, 625-633. Hughes, J. R. (1995). Combining behavioral therapy and pharmacotherapy for smoking cessation: An update. *NIDA Research Monograph*, 150, 92-109. Gulliver, S. B., Kamholz, B. W., & Helstron, A. W. (2006). Smoking cessation and alcohol abstinence: What do the data tell us? *Alcohol Research & Health*, 29(3), 208-212. York, J. L., & Hirsch, J. A. (1995). Drinking patterns and health status in smoking and nonsmoking alcoholics. *Alcoholism: Clinical and Experimental Research*, 19, 666-673.
- ¹⁹ Hays, J.T., Schroeder, D.R., Offord, K.P., Croghan, I.T., Patten, C.A., Hurt, R.D.,...Fiore, M. C. (1999). Response to nicotine dependence treatment in smokers with current and past alcohol problems. *Annals of Behavioral Medicine*, 21(3), 244-250.
- ²⁰ Ong, M. K., Zhou, Q., & Sung, H-Y. (2011). Primary care providers advising smokers to quit: Comparing effectiveness between those with and without alcohol, drug, or mental disorders. *Nicotine & Tobacco Research*, August 23, DOI: [10.1093/ntr/ntr167](https://doi.org/10.1093/ntr/ntr167)
- ²¹ Citations from Knudsen, H. K., & Studts, J. L. (2010). The implementation of tobacco-related brief interventions in substance abuse treatment: A national study of counselors. *Journal of Substance Abuse Treatment*, 38, 212-219: Kalman, D., Hayes, K., Colby, S. M., Easton, C. A., Rohsenow, D. J., & Monti, P. M. (2001). Concurrent

- versus delayed smoking cessation treatment for persons in early alcohol recovery. A pilot study. *Journal of Substance Abuse Treatment*, 20, 233-238. Kalman, D., Kim, S., DiGirolam, G., Smelson, D., & Ziedonis, Z. (2010). Addressing tobacco use disorder in smokers in early remission from alcohol dependence. *Clinical Psychology Review*, 30(1), 12-24. McCarthy, W. J., Collins, C., & Hser, Y.I. (2002). Does smoking cessation effect drug abuse treatment? *Journal of Drug Issues*, 32, 61-80. Richter, K. P., Choi, W. S., McCool, R. M., Harris, K. J., & Ahluwalia, J. S. (2002). A population-based study of cigarette smoking among illicit drug users in the United States. *Addiction*, 97, 861-869. Williams, J. M., & Ziedonis, D. (2004). Addressing tobacco among individuals with a mental illness of an addiction. *Addictive Behaviors*, 29, 1067-1083.
- ²² Burling, T.A., Bulring, A.S., & Latini, D. (2001). A controlled smoking cessation trial for substance dependent inpatients. *Journal of Consulting and Clinical Psychology*, 69, 295-304.
- ²³ Hser, Y.I, McCarthy, W.J., & Anglin, M.D. (1994). Tobacco use as a distal predictor of mortality among long-term narcotic addicts. *Prevention Medicine*, 23, 61-69. Hurt, R.D., Offord, K.P., Vroghanm I.T., Gomez-Dahl, L., Kottke, T.E. Morse, R.M., & Melton, J. (1996). Mortality following inpatient addictions treatment. *Journal of the American Medical Association*, 274(14), 1097-1103.
- ²⁴ For biographical information, see White, W. (1998). *Slaying the dragon: The history of addiction treatment and recovery in America*. Bloomington, IL: Chestnut Health Systems.
- ²⁵ Citations from Knudsen, H. K., & Studts, J. L. (2010). The implementation of tobacco-related brief interventions in substance abuse treatment: A national study of counselors. *Journal of Substance Abuse Treatment*, 38, 212-219; Lemon, S.C., Friedmann, P.D., & Stein, M.D. (2003). The impact of smoking cessation on drug abuse treatment outcome. *Addictive Behaviors*, 28, 1323-1331. McCarthy, W.J., Collins, C., & Hser, Y.I. (2002). Does smoking cessation effect drug abuse treatment? *Journal of Drug Issues*, 32, 61-80. Satre, D.D., Kohn, C.S., & Weisner, C. (2007). Cigarette smoking and long-term alcohol and drug treatment outcomes: A telephone follow-up at five years. *American Journal on Addictions*, 16, 32-37.
- ²⁶ Cooney, J.L., Cooney, N.L., Pilkey, D.T., Krnazler, H.R., & Onken, C.A. (2003). Effects of nicotine deprivation on urges to drink and smoke in alcoholic smokers. *Addiction*, 98, 913-921.
- ²⁷ Sobell, M.B., Sobell, L.C., & Kozlowski, L.T. (1995). Dual recoveries from alcohol and smoking problems. In J.B. Fertig, & J.A. Allen (Eds.), *Alcohol and tobacco: From basic science to clinical practice* (NIAAA Research Monograph No. 30, pp. 207-224). Rockville, MD: NIAAA. Taylor, B.J., Graham, J.W., Cumsille, P., & Hansen, W. B. (2000). Modeling prevention program effects on growth of substance use: Analysis of five years of data from the Adolescent Alcohol Prevention Trial. *Prevention Science*, 1, 183-197.
- ²⁸ Kalman, D., Kim, S., DiGirolam, G., Smelson, D., & Ziedonis, Z. (2010). Addressing tobacco use disorder in smokers in early remission from alcohol dependence. *Clinical Psychology Review*, 30(1), 12-24. Prochaska, J.J., Delucchi, K., & Hall, S.M. (2004). A meta-analysis of smoking cessation interventions with individuals in substance abuse treatment or recovery. *Journal of Consulting and Clinical Psychology*, 72(6), 1144-1156.
- ²⁹ Kohn, C.S., Tsoh, J.Y., & Weisner, C.M. (2003). Changes in smoking status among substance abusers: Baseline characteristics and abstinence from alcohol and drugs at 12-month follow-up. *Drug and Alcohol Dependence*, 69, 61-71.
- ³⁰ Fiore, M.C., Jaen, C.R., Baker, T.B., Bailey, W.C., Benowitz, N.L., Curry, S.J.,...Wewers, M. E. (2008). *Treating tobacco use and dependence: 2008 update. Clinical practice guidelines*. Rockville, MD: U.S. Department of Health and Human Services. Public Health Service. Silagy, C., Lancaster, T., Stead, L., Mant, D., & Fowler, G. (2003). Nicotine replacement therapy for smoking cessation (Cochrane Review). In *The Cochrane Library*, Issue, 1.
- ³¹ Prochaska, J.J., Delucchi, K., & Hall, S.M. (2004). A meta-analysis of smoking cessation interventions with individuals in substance abuse treatment or recovery. *Journal of Consulting and Clinical Psychology*, 72(6), 1144-1156.
- ³² Fiore, M.C., Jaen, C.R., Baker, T.B., Bailey, W.C., Benowitz, N.L., Curry, S.J.,...Wewers, M. E. (2008). *Treating tobacco use and dependence: 2008 update. Clinical practice guidelines*. Rockville, MD: U.S. Department of Health and Human Services. Public Health Service.
- ³³ Ong, M. K., Zhou, Q., & Sung, H-Y. (2011). Primary care providers advising smokers to quit: Comparing effectiveness between those with and without alcohol, drug, or mental disorders. *Nicotine & Tobacco Research*, August 23, DOI: [10.1093/ntr/ntr167](https://doi.org/10.1093/ntr/ntr167)
- ³⁴ Monti, P.M., Rohsenow, D.J., Colby, S.M., & Abrams, D.B. (1995). Smoking among alcoholics during and after treatment: Implications for models, treatment strategies and policy. In J.B. Fertig & J.P. Allen (Eds.), *Alcohol and tobacco from basic science to clinical practice* (NIAAA Research Monograph, Vol 30, p. 187-206). Bethesda, MD: National Institutes of Health.

-
- ³⁵ Karan, L.D. (1993). Towards a broader view of recovery. *Journal of Substance Abuse Treatment*, *10*(2), 101-105. McLellan, A. T. (2010). What is recovery? Revisiting the Betty Ford Institute Consensus Panel definition. *Journal of Substance Abuse Treatment*, *38*, 200-201. doi:10.1016/j.jsat.2009.11.002 White, W. L. (2007). Addiction recovery: Its definition and conceptual boundaries. *Journal of Substance Abuse Treatment*, *33*, 229-241. doi:10.1016/j.jsat.2007.04.015
- ³⁶ *Smoking Cessation Fact Sheet*. (2010). Washington D.C.: American Legacy Foundation. Citations available online at <http://www.legacyforhealth.org/>.
- ³⁷ Kalman, D., Kim, S., DiGirolam, G. Smelson, D., & Ziedonis, Z. (2010). Addressing tobacco use disorder in smokers in early remission from alcohol dependence. *Clinical Psychology Review*, *30*(1), 12-24.
- ³⁸ CDC. (2009). Cigarette smoking among adults and trends in smoking cessation—United States, 2008. *MMWR*, *58*(44), 1227-1232.
- ³⁹ Hughes, J.R., Callas, P.W., & High Dose Study Group. (2003). Past alcohol problems do not predict worse smoking cessation outcomes. *Drug and Alcohol Dependence*, *71*(3), 269-273. Hughes, J.R. & Kalman, D. (2006). Do smokers with alcohol problems have more difficulty quitting? *Drug and Alcohol Dependence*, *82*(2), 91-102.
- ⁴⁰ Falba, T. A., & Sindelar, J. L. (2008). Spousal concordance in health behavior change. *Health Services Research*, *43*(1), 96 -116.
- ⁴¹ Chapman, S, & MacKenzie, R. (2010). *The global research neglect of unassisted smoking cessation: Causes and consequences*. PLoS Med *7*(2): e1000216. doi:10.1371/journal.pmed.1000216
- ⁴² Zhu, S-H., Melcer, T., Sun, J., Rosbrook, B. & Pierce, J.P. (2000). Smoking cessation with and without assistance: A population-based analysis. *American Journal of Preventive Medicine*, *18*(4), 305-311.
- ⁴³ Martin, M.E., Calfas, K.J., Patten, C.A., Polarek, M., Hofstetter, C.R., Noto, J., & Beach, D. (1997). Prospective evaluation of three smoking interventions in 205 recovering alcoholics: One-year results of Project SCRAP-tobacco. *Journal of Consulting and Clinical Psychology*, *65*, 190-194.
- ⁴⁴ Hughes, J.R., & Kalman, D. (2006). Do smokers with alcohol problems have more difficulty quitting? *Drug and Alcohol Dependence*, *82*(2), 91-102.